

McLean Health Center Facility Quality Measures/Indicators

High Risk/Management Priority (Lower Percentages are Better)

Quality Measure/Indicator		2006 Q2	2006 Q3	2006 Q4	2007 Q1	2007 Q2	2007 Q3	2007 Q4	2008 Q1
Incidence of new fractures	McLean	2.8	4.0	1.3	1.6	1.1	1.2	2.1	1.2
	CT Rate	1.6	1.7	1.7	1.7	1.8	1.8	1.9	1.9
Prevalence of falls	McLean	8.0	8.3	8.7	6.2	9.4	9.9	10.4	13.2
	CT Rate	12.0	11.7	11.8	11.8	11.9	11.9	11.9	12.9
Prevalence of fecal impaction	McLean	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
	CT Rate	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Prevalence of dehydration	McLean	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
	CT Rate	0.4	0.3	0.5	0.4	0.4	0.4	0.4	0.4
High risk residents with pressure ulcers	McLean	14.3	9.1	6.6	11.9	8.7	6.9	8.4	3.9
	CT Rate	14.4	13.5	14.1	14.1	13.7	13.7	13.5	13.6
Low risk residents with pressure ulcers	McLean	0.0	3.1	0.0	0.0	0.0	0.0	0.0	0.0
	CT Rate	2.7	2.3	2.7	2.6	2.5	2.5	2.7	2.8

McLean Health Center Facility Quality Measures/Indicators

Other Important Indicators for State and Federal Compliance (Lower Percentages are Better)

Quality Measure/Indicator		2006 Q2	2006 Q3	2006 Q4	2007 Q1	2007 Q2	2007 Q3	2007 Q3	2008 Q1
Residents with a urinary tract infection	McLean	7.1	7.5	4.3	4.1	5.1	3.0	7.8	11.0
	CT Rate	8.1	8.2	8.1	8.1	8.2	8.2	8.3	8.3
Residents who lose too much weight	McLean	5.3	6.9	5.9	8.5	10.9	9.9	8.9	6.1
	CT Rate	10.0	9.6	9.7	9.7	9.5	9.5	9.6	9.7
Prevalence of antipsychotic use, in the absence of psychotic or related conditions: Overall	McLean	25.0	26.0	23.4	25.0	24.5	23.8	25.8	22.2
	CT Rate	26.6	26.9	26.8	26.7	26.9	26.9	26.6	26.4
Prevalence of antipsychotic use, in the absence of psychotic or related conditions: High risk	McLean	42.1	30.0	50.0	46.7	47.4	41.7	63.6	50.0
	CT Rate	55.6	56.6	55.8	55.9	55.4	55.4	55.0	55.2

Quality Measure/Indicator		2006 Q2	2006 Q3	2006 Q4	2007 Q1	2007 Q2	2007 Q3	2007 Q4	2008 Q1
Prevalence of antipsychotic use, in the absence of psychotic or related conditions: Low risk	McLean	20.3	25.6	18.5	19.7	18.7	20.6	20.7	20.0
	CT Rate	23.5	23.9	23.6	23.6	24.1	24.1	23.9	23.7
Residents who were physically restrained	McLean	4.5	3.3	4.3	5.2	2.6	0.0	0.0	0.0
	CT Rate	4.6	4.2	4.8	4.6	4.2	4.2	3.9	3.9
Residents who have become more depressed or anxious	McLean	5.9	19.1	25.3	30.9*	26.9	15.4	9.2	7.5
	CT Rate	13.9	13.9	13.7	13.8	13.9	13.9	14.1	14.1
Prevalence of behavior symptoms affecting others: Overall	McLean	24.1	12.5	19.6	19.6	19.7	12.9	12.2	7.7
	CT Rate	11.8	11.7	12.0	11.9	11.4	11.4	11.1	11.1
Prevalence of behavior symptoms affecting others: High risk	McLean	27.1	13.7	20.0	22.2	21.4	15.7	13.4	8.1
	CT Rate	13.7	13.6	13.9	13.8	13.2	13.2	12.9	12.9
Prevalence of behavior symptoms affecting others: Low risk	McLean	6.3	6.3	18.2	7.7	6.3	0.0	0.0	0.0
	CT Rate	4.9	4.6	5.1	5.0	4.7	4.7	4.3	4.3
Prevalence of symptoms of depression without antidepressant therapy	McLean	1.8	2.5	3.3	7.2	11.1	4.0	1.7	1.1
	CT Rate	2.9	2.9	3.0	3.0	3.0	3.0	3.0	3.0
Use of nine or more different medications	McLean	68.8	72.5	67.4	71.1	71.8	73.3	75.7	81.3
	CT Rate	63.0	63.8	62.4	62.8	64.1	64.1	65.0	65.1
Incidence of cognitive impairment	McLean	0.0	5.3	12.5	0.0	0.0	6.3	10.0	5.6
	CT Rate	13.0	12.7	12.6	12.6	13.0	13.0	13.2	13.3
Low risk residents who lost control of their bowels or bladder	McLean	67.1	57.8	61.0	66.2	67.1	75.7	73.5	75.4
	CT Rate	45.4	46.1	45.4	45.5	46.2	46.2	46.9	47.1

Quality Measure/Indicator		2006 Q2	2006 Q3	2006 Q4	2007 Q1	2007 Q2	2007 Q3	2007 Q4	2008 Q1
Residents who have/had a catheter inserted and left in their bladder	McLean	13.4	13.3	8.7	8.0	9.4	7.9	9.2	10.8
	CT Rate	5.8	5.5	5.7	5.8	5.8	5.8	5.7	5.7
Prevalence of occasional or frequent bladder or bowel incontinence without a toileting plan	McLean	8.0	7.7	15.9	10.4	13.8	18.9	16.4	25.0
	CT Rate	37.6	39.4	36.7	37.3	39.8	39.8	40.6	40.6
Prevalence of tube feeding	McLean	2.7	2.5	2.2	1.0	1.7	2.0	1.7	3.3
	CT Rate	3.6	3.6	3.6	3.6	3.6	3.6	3.5	3.5
Residents who have moderate to severe pain	McLean	5.4	7.5	9.8	6.0	4.0	5.0	5.1	1.6
	CT Rate	7.2	7.0	7.3	4.7	4.6	4.6	4.4	4.3
Residents whose need for help with daily activities has increased	McLean	11.6	11.8	16.7	24.2	16.1	11.9	7.5	16.7
	CT Rate	19.3	18.0	18.5	18.6	18.8	18.8	19.1	19.3
Residents who spend most of their time in bed or in a chair	McLean	2.7	1.7	1.1	0.0	0.0	0.0	0.0	0.0
	CT Rate	2.5	2.4	2.5	2.4	2.6	2.6	2.7	2.8
Residents whose ability to move in and around their room got worse	McLean	15.2	13.2	15.6	31.9	23.2	25.5	9.7	9.0
	CT Rate	17.9	17.8	17.6	16.9	17.3	17.3	17.6	17.7
Incidence of decline in range of motion	McLean	6.5	5.7	1.2	4.3	5.1	11.6	5.0	11.2
	CT Rate	6.7	6.6	6.6	6.6	6.7	6.7	6.7	6.7
Prevalence of antianxiety/hypnotic use	McLean	19.3	24.5	26.9	26.7	28.3	28.6	26.5	31.7
	CT Rate	21.2	21.3	21.3	21.3	21.7	21.7	22.0	22.0
Prevalence of hypnotic use more than two times in last week	McLean	3.6	5.8	5.4	7.2	6.0	8.9	7.8	11.0
	CT Rate	6.1	6.2	6.1	6.2	6.2	6.2	6.2	6.2
Prevalence of little or no activity	McLean	0.9	0.8	0.0	0.0	0.0	0.0	0.0	0.0
	CT Rate	3.1	3.0	3.1	3.0	3.0	3.0	2.9	2.9